

Forecasting the VIX in the midst of COVID-19

Rutger Lit^{(a,b)*}

^(a)Time Series Lab

^(b)Vrije Universiteit Amsterdam, The Netherlands

April 18, 2020

Abstract

We study the behavior of the Volatility Index (VIX) time series in the period leading up to the COVID-19 outbreak. Time-varying location/scale models are used to extract a range of unobserved components from the VIX time series. The time-varying unobserved components are driven by the score of the predictive density. These so called score-driven models have proven to be powerful in extracting unobserved components like autoregressive processes and seasonal patterns. A range of model specifications is used to forecast the VIX in the COVID-19 period that spans the first quarter of 2020. Explanatory variables are used to improve in-sample model fit and out-of-sample forecast accuracy. All model computations are carried out with the *Time Series Lab* software package.

Key words: VIX, COVID-19, Time Series Lab, Unobserved components, Time Series, Forecasting, Score-driven models

1 Introduction

The VIX, also called fear index or fear gauge, is a measure of the overall market sentiment and reflects the investors' risk appetite. The VIX is calculated by the Chicago Board Options Exchange (CBOE) and estimates implied volatility by aggregating the weighted prices of S&P 500 put and call options over a wide range of strike prices. A detailed description of the calculation of the VIX is given by CBOE in their white paper, see [CBOE \(2019\)](#). The VIX is quoted in percentage points and represents the expected annualized change in the S&P 500 index over the next 30-day time period within a one standard deviation confidence interval. However, VIX today more often than not overstates the level of actual volatility experienced in the next 30 days, see [Edwards and Preston \(2017a\)](#) and [Edwards and Preston \(2017b\)](#). Two explanations have been given, the first one stems from a behavioral finance perspective. Risk averse investors are willing to pay a premium to hedge against risk. The second one comes from an option pricing perspective. Implied volatility takes into account rare events that occur in the (often non-Gaussian) tails of

*The author is grateful for comments from Paolo Gorgi, Siem Jan Koopman, and Binte Groen. All errors are his own.

the distribution, while realized volatility will only include rare events if they have occurred in the period for which the realized volatility is calculated. Since on average rare events will only occur with a small frequency, realized volatility tends to underestimate the potential for large losses most of the time.

In this paper we present our study into the behavior of the Volatility Index (VIX) time series and forecast daily closing levels of the VIX for the first quarter of 2020. This time period was chosen deliberately because in March 2020 the VIX fluctuated as never seen before due to the COVID-19 outbreak. We focused on time-varying location/scale models with the aim of identifying the model with the most accurate forecast of the VIX in turbulent times. With accurate forecasts of implied volatility, market participants could potentially form profitable option trading strategies, something which can have implications about the efficiency of the option markets as well. The VIX data and additional explanatory variables of this research have a daily frequency, which means that we calculate forecasts for day $t + 1$ with information up to and including time t . The author of this paper is fully aware that in today's high-frequency markets and 24h economy, forecasting one day ahead for a liquid product is a long forecast horizon and by the time the forecast period is reached, new information has already become available that could potentially improve forecasts. However, if statistically significant conclusions can be drawn from forecasts one day ahead, it can assist us in decision making processes at the current day.

We modelled the VIX time series with a wide range of probability distributions, dynamic components, and explanatory variables and compared these models on in-sample model fit and out-of-sample forecast accuracy. All model calculations in this paper were carried out with the software package [Time Series Lab - Score Edition](https://timeserieslab.com) which is available from <https://timeserieslab.com> for free and can therefore be easily replicated. Screen captures of the software are presented in the online supplement to highlight the chosen settings in each step of the modelling process.

For more information on modelling and forecasting of implied volatility, we refer to [Fernandes et al. \(2014\)](#), [Koopman et al. \(2005\)](#), [Psaradellis and Sermpinis \(2016\)](#), [Konstantinidi et al. \(2008\)](#), [Blair et al. \(2001\)](#), [Blair et al. \(2010\)](#), [Harvey and Whaley \(1992\)](#) and the references in these papers.

2 The model

The VIX time series is assumed to have observations coming from a probability distribution $p(y_t|\mu_t, \sigma_t)$ with time-varying location μ_t and scale σ_t for $t = 1, \dots, T$ where T is the length of the time series. The mechanism to update the location and scale over time is the scaled score of the likelihood function. These so called score-driven models, or Generalized Autoregressive Score models, were proposed in their full generality by [Creal et al. \(2013\)](#) and for time-varying location/scale volatility models by [Harvey \(2013\)](#). The score-driven approach provides a unified and consistent framework for introducing time-varying parameters in a wide class of nonlinear models. Score-driven models encompass several well-known models like the GARCH model of [Engle \(1982\)](#) and the ACD model of [Engle and Russell \(1998\)](#). [Time Series Lab - Score Edition](#) specializes in models with unobserved components driven by the score.

Let the unobserved time-varying parameter vector α_t be updated over time using the following

updating function

$$\alpha_{t+1} = \omega + \sum_{i=1}^q A_i s_{t-i+1} + \sum_{j=1}^p B_j \alpha_{t-j+1},$$

where ω is a vector of constants, A and B are fixed coefficient matrices and s_t is the scaled score function which is the driving force behind the updating equation. The unknown coefficients ω , A and B depend on the static parameter vector ψ which is estimated by the method of Maximum Likelihood. The definition of s_t is

$$s_t = S_t \cdot \nabla_t, \quad \nabla_t = \frac{\partial \log p(y_t | \alpha_t, \mathcal{F}_{t-1}; \psi)}{\partial \alpha_t},$$

for $t = 1, \dots, T$, and where ∇_t is the score vector of the density $p(y_t | \alpha_t, \mathcal{F}_{t-1}; \psi)$. The information set \mathcal{F}_{t-1} usually consists of lagged variables of α_t and y_t but can contain exogenous variables as well. The connection between the distribution parameters μ_t and σ_t is made by the link function $f(\cdot)$ in the following way

$$\mu_t = f(Z_t^\mu \alpha_t), \quad \sigma_t = f(Z_t^\sigma \alpha_t),$$

where at time t , the selection matrices Z_t^μ and Z_t^σ select components from the unobserved parameter vector α_t .

In *Time Series Lab*, a wide range of unobserved components can be included in α_t . Among these are non-stationary components like Trend and Seasonal, and stationary components like Autoregressive processes of order p . The score-driven framework easily takes explanatory variables into account as well, something which we will use for the modelling of the VIX time series. Furthermore, *Time Series Lab* allows the user to choose from several probability distributions $p(y_t | \mu_t, \sigma_t)$.

3 Data description

The daily VIX time series was downloaded from Yahoo finance (<https://finance.yahoo.com>) and descriptive statistics are presented in Table 1. Note that the downloaded raw data has weekends and banking holidays removed so that no missing values are present in the data. To be able to model a possible weekly, or biweekly pattern we added missing values for the banking holidays so that each 5 consecutive data points constitutes exactly 1 week. The VIX time series is presented in Figure 1 for the full sample and the first quarter of 2020. Our approach is to estimate model parameters from period 1 and use these to forecast period 2. This poses several challenges since both periods differ strongly when it comes to their statistical characteristics. For example, mean and standard deviation are much higher in period 2. Also, the data is skewed which is somewhat alleviated after taking logs from the series but not completely. Furthermore, kurtosis switches from excess kurtosis (> 3) in period 1 to leptokurtic (< 3) in period 2. Most noteworthy is the strong rejection of a unit root in period 1 and the strong acceptance of a unit root in period 2.

The following lag-1 explanatory variables were used in an attempt to increase forecast accuracy: the κ -day S&P 500 continuously compounded positive returns and negative returns for $\kappa = 1, 5, 10, 20$. The κ -day continuously compounded return of WTI Crude oil price. The S&P 500 volume first difference of logs. The first difference of the logarithm of the foreign

exchange value of the US dollar against the Euro, British pound, Japanese yen, and Bitcoin. The spread between 10-Year and 3-month Treasury Constant Maturities (credit spread). The difference between the three-month Treasury bill and the three-month LIBOR based in US dollars (TED spread) and dummy variables for the beginning and end of the trading month.

Explanatory variables were downloaded from Yahoo finance and the Federal Reserve Economic Data base (<https://fred.stlouisfed.org>).

4 Results

The ACF of the VIX time series shows a slowly decaying pattern while the PACF shows 2 significant lags. We therefore started our analysis with a benchmark model consisting of a time-varying location that consists of an AR2 process and a constant scale parameter. The most complex model has a location parameter that is decomposed into an AR2 process, an AR1 process, a weekday seasonal specification, explanatory variables, and a scale parameter consisting of an AR1 process. All components, except the explanatory variables component, are driven by the score as described in Section (2). In-sample model fit and out-of-sample forecast accuracy is reported and compared among other models including simple benchmark models in Table 2. The losses were evaluated by the Diebold and Mariano (1995) statistic to test for equal predictive accuracy. A higher (in-sample) log likelihood value results in almost all cases in lower in-sample loss, however, this is not true for out-of-sample forecast accuracy. The more complex and better fitting in-sample models are often not better at forecasting as indicated by the higher out-of-sample losses. This can be explained by overfitting of the time series, a phenomenon where simple time series models outperform more complex models when it comes to forecasting. We discuss estimation results per category.

Distribution

Looking at point forecasts (RMSE, MAE, MAPE), the Gaussian distribution outperforms the Student t and Generalized Error Distribution (GED) in several cases and is never significantly worse.

This strong performance on point forecasting is in contrast to the performance when the full distribution is taken into account, e.g. better tail fitting behavior. The in-sample log likelihood value is much higher for the Student t distribution, a clear sign of fatter tails compared to the Gaussian distribution. The parameter estimates are presented in Table 3 from which we see that the degrees of freedom parameter is estimated at 7.40 indicating fatter tails than the Gaussian distribution. The log loss function, which is defined as minus the log of the predictive density is also lower for the Student t models.

These results might be surprising at first but are actually in line with theory, since the Gaussian distribution minimizes the squared errors. It is for the practitioner to decide which is valued higher, point forecasts or density forecasts.

Autoregressive components

Time Series Lab allows the inclusion of several unobserved components. The extracted components for location and scale are presented in Figure 2 and 3. We note that the signal for location

is decomposed into a persistent AR2 processes ($\phi_1 + \phi_2 = 0.996$) and an AR1 process that takes the short(er) run shocks into account albeit relatively persistent as well with ($\phi = 0.863$). The sum of two autoregressive components can be seen as a long memory process, see (Harvey, 2013, p91) for long memory in location/scale models and Fernandes et al. (2014) for long memory in VIX modelling.

Seasonal

A small but significant weekday seasonal is extracted from the VIX time series. Table 2 shows the parameters of the seasonal components. First, the seasonal is not time-varying since the updating parameter (seasonal κ) was estimated close to zero and subsequently fixed at zero during re-estimation. Second, there is a weekday effect in which the VIX is on average slightly higher on Monday and Tuesday and lower on Friday. The parameters for Wednesday and Thursday are not significantly different from zero. Note that the seasonal components must sum to zero for identification and therefore the Friday parameter can be deduced from the parameters belonging to the rest of the week. We refer to Figure 2 for a plot of the weekday pattern.

Explanatory variables

After optimizing the model with the full set of explanatory variables, corresponding standard errors were calculated. The least significant explanatory variable was removed and the process of estimating and calculating standard errors was repeated until a significant set of explanatory variables remained. The final model consists of the explanatory variables S&P negative return lag-1, WTI Close return lag-1 (oil price), and a dummy for the beginning and end of the trading month. We refer to Table 2 for the parameter estimates and standard errors of the explanatory variables. The negative sign in front of the coefficients corresponding to S&P500 negative returns lag1 and WTI closing price lag1 show an inverse relationship with the VIX.

Model extensions

The inclusion of multiple lags of the autoregressive processes was tried for both location and scale but without improvement. Furthermore, a seasonal pattern could not be extracted from the scale parameter. Finally, multiple lags of the score were included in the model as well without improvement on model fit or forecast-accuracy.

5 Conclusion

We used the time series package [Time Series Lab - Score Edition](#) to extract signal from the VIX time series. We modelled the time-varying location and scale of the model for several probability distributions and combinations of unobserved components. Especially the time-varying location parameter was a rich source of information since two autoregressive processes and a weekday seasonal pattern was extracted.

The Gaussian distribution outperformed the Student t and Generalized Error Distribution (GED) when it comes to point forecasts (RMSE, MAE, MAPE). This strong performance on point forecasting is in contrast to the performance when the full distribution is taken into account.

These results are in line with theory since the Gaussian distribution minimizes the squared errors. We found long memory in the VIX time series due to the 2 persistent AR process that were extracted. A small but significant weekday seasonal is extracted from the VIX time series as well. There is a weekday effect in which the VIX is on average slightly higher on Monday and Tuesday and lower on Friday. The final model consists of the explanatory variables S&P negative return lag-1, WTI Close return lag-1, and a dummy for the beginning and end of the trading month. The negative sign in front of the coefficients corresponding to S&P500 negative returns lag1 and WTI closing price lag1 show an inverse relationship with the VIX.

References

- Blair, B. J., S.-H. Poon, and S. J. Taylor (2001). Modelling s&p 100 volatility: The information content of stock returns. *Journal of banking & finance* 25(9), 1665–1679.
- Blair, B. J., S.-H. Poon, and S. J. Taylor (2010). Forecasting s&p 100 volatility: the incremental information content of implied volatilities and high-frequency index returns. In *Handbook of Quantitative Finance and Risk Management*, pp. 1333–1344. Springer.
- CBOE (2019). Vix white paper cboe volatility index[®]. <http://www.cboe.com/micro/vix/vixwhite.pdf>.
- Creal, D. D., S. J. Koopman, and A. Lucas (2013). Generalized autoregressive score models with applications. *Journal of Applied Econometrics* 28(5), 777–795.
- Diebold, F. X. and R. S. Mariano (1995). Comparing predictive accuracy. *Journal of Business and Economic Statistics* 13, 253–265.
- Edwards, T. and H. Preston (2017a). A practitioners guide to reading vix[®].
- Edwards, T. and H. Preston (2017b). Reading vix[®]: Does vix predict future volatility?
- Engle, R. F. (1982). Autoregressive conditional heteroscedasticity with estimates of the variance of united kingdom inflation. *Econometrica*, 987–1007.
- Engle, R. F. and J. R. Russell (1998). Autoregressive conditional duration: a new model for irregularly spaced transaction data. *Econometrica*, 1127–1162.
- Fernandes, M., M. C. Medeiros, and M. Scharth (2014). Modeling and predicting the cboe market volatility index. *Journal of Banking & Finance* 40, 1–10.
- Harvey, A. C. (2013). *Dynamic Models for Volatility and Heavy Tails: With Applications to Financial and Economic Time Series*, Volume 52. Cambridge: Cambridge University Press.
- Harvey, C. R. and R. E. Whaley (1992). Market volatility prediction and the efficiency of the s & p 100 index option market. *Journal of Financial Economics* 31(1), 43–73.
- Konstantinidi, E., G. Skiadopoulos, and E. Tzagkaraki (2008). Can the evolution of implied volatility be forecasted? evidence from european and us implied volatility indices. *Journal of Banking & Finance* 32(11), 2401–2411.


Koopman, S. J., B. Jungbacker, and E. Hol (2005). Forecasting daily variability of the s&p 100 stock index using historical, realised and implied volatility measurements. *Journal of Empirical Finance* 12(3), 445–475.

Lit, R., S. J. Koopman, and A. C. Harvey (2019-2020). Time Series Lab - Score Edition. <https://timeserieslab.com>.

Psaradellis, I. and G. Sermpinis (2016). Modelling and trading the us implied volatility indices. evidence from the vix, vxn and vxd indices. *International Journal of Forecasting* 32(4), 1268–1283.

Tables and Figures

Figure 1
VIX closing levels



The figure shows closing levels of the VIX. Top panel: full sample. Bottom panel: COVID-19 outbreak period. The top panel clearly shows the sharp rise of the VIX in a very short time period at the end of the sample. Equal levels were reached during the financial crisis however the period leading up to the financial crisis was more turbulent than the period leading up to the COVID-19 jump. The bottom panel shows that the VIX calmly started the beginning of 2020 and then abruptly rose.

Table 1
Descriptive statistics of the VIX time series

The table reports descriptive statistics of the VIX time series for the closing price and log closing price. Period 1 ranges from 05-01-2004 to 31-12-2019. Period 2 spans the first quarter of 2020 in which the COVID-19 outbreak took place (01-01-2020 to 31-03-2020). The p-values of the Jarque-Bera test for normality and the Augmented Dickey-Fuller (ADF) test for unit root are reported. The number of lags in the ADF test are selected using the Bayesian information criterion.

Characteristic	Closing price			Log closing price		
	Period 1	Period 2	Full sample	Period 1	Period 2	Full sample
Mean	18.22	31.22	18.42	2.83	3.21	2.83
Standard dev.	8.58	22.05	9.08	0.36	0.67	0.37
Median	15.60	16.73	15.61	2.75	2.82	2.75
Minimum	9.14	12.10	9.14	2.21	2.49	2.21
Maximum	80.86	82.69	82.69	4.39	4.42	4.42
Kurtosis	13.41	2.27	13.74	4.57	1.59	4.73
Skewness	2.75	0.88	2.84	1.17	0.50	1.23
T	4172	66	4238	4172	66	4238
Missings	146	4	150	146	4	150
Jarque-Bera	0.0000	0.0068	0.0000	0.0000	0.0164	0.0000
ADF	0.0002	0.9502	0.0019	0.0000	0.9516	0.0000

Table 2

In-sample and out-of-sample results for a range of model specifications

The table reports in-sample model fit and out-of-sample forecast results for a wide range of model specifications. All models are estimated with [Time Series Lab - Score Edition](#). All dynamic components are driven by the score. The lowest loss per category is given a blue shade. The [Diebold and Mariano \(1995\)](#) statistic is used to assess forecast accuracy among competing models. The second Student t model with Location: $\exp(\text{AR2})$ and Scale: $\exp(\text{AR1})$ is set as benchmark model. All out-of-sample losses with a dagger sign † next to them perform significantly worse compared to the benchmark model.

Model description	In-sample				Out-of-sample			
	Log L	RMSE	MAE	MAPE	RMSE	MAE	MAPE	Log loss
Distribution: Gaussian								
Location: $\exp(\text{AR2})$ Scale: constant	-7895.50	1.72	1.02	5.18	5.91	3.37	8.64	7.36†
Location: $\exp(\text{AR2})$ Scale: $\exp(\text{AR1})$	-6687.61	1.74	1.01	5.13	6.05	3.40	8.63	2.91
Location: $\exp(\text{AR2} + \text{AR1})$ Scale: $\exp(\text{AR1})$	-6649.01	1.74	1.00	5.04	6.49	3.44	8.55	2.94
Location: $\exp(\text{AR2} + \text{AR1} + \text{Seas}(5))$ Scale: $\exp(\text{AR1})$	-6589.67	1.74	0.99	4.97	6.47	3.53	8.80	2.91
Location: $\exp(\text{AR2} + \text{AR1} + \text{Seas}(5) + X\beta)$ Scale: $\exp(\text{AR1})$	-6536.45	1.69	0.98	4.93	6.27	3.68	9.09	2.90
Distribution: Student t								
Location: $\exp(\text{AR2})$ Scale: constant	-7293.05	2.05	1.11	5.49	18.14†	11.09†	21.89†	7.15†
Location: $\exp(\text{AR2})$ Scale: $\exp(\text{AR1})$	-6306.96	1.80	1.04	5.19	5.96	3.33	8.86	2.61
Location: $\exp(\text{AR2} + \text{AR1})$ Scale: $\exp(\text{AR1})$	-6270.03	1.83	1.03	5.13	6.70†	3.64	9.34	2.65
Location: $\exp(\text{AR2} + \text{AR1} + \text{Seas}(5))$ Scale: $\exp(\text{AR1})$	-6209.58	1.83	1.03	5.06	6.64†	3.68	9.49	2.66
Location: $\exp(\text{AR2} + \text{AR1} + \text{Seas}(5) + X\beta)$ Scale: $\exp(\text{AR1})$	-6162.03	1.80	1.01	5.00	7.16†	4.09†	10.08†	2.69
Distribution: GED								
Location: $\exp(\text{AR2})$ Scale: constant	-7485.44	1.82	1.08	5.50	7.11†	3.99†	10.20†	6.33†
Location: $\exp(\text{AR2})$ Scale: $\exp(\text{AR1})$	-6464.19	1.81	1.04	5.24	6.00	3.26	8.60	2.71
Location: $\exp(\text{AR2} + \text{AR1})$ Scale: $\exp(\text{AR1})$	-6428.11	1.85	1.04	5.19	6.57	3.43	8.83	2.74
Location: $\exp(\text{AR2} + \text{AR1} + \text{Seas}(5))$ Scale: $\exp(\text{AR1})$	-6367.55	1.84	1.03	5.11	6.54	3.42	8.83	2.74
Location: $\exp(\text{AR2} + \text{AR1} + \text{Seas}(5) + X\beta)$ Scale: $\exp(\text{AR1})$	-6317.81	1.78	1.01	5.04	7.11†	3.95†	9.64	2.78

Table 3
Parameter estimates for the Student t model

The table reports parameter estimates obtained from maximizing the likelihood function of the most complex Student t model with specification Location: $\exp(\text{AR2}+\text{AR1}+\text{Seas}(5)+X\beta)$ and Scale: $\exp(\text{AR1})$. P-values should be taken with caution. They are directly interpretable for regression coefficients but, due to boundary issues, can give misleading results for parameters corresponding to dynamic components or variances.

Parameter type	Value	Sig.Lvl	Asymp.SE	p-value	-1.96 SE	+1.96 SE
Log location						
AR2 ω	0.0091	***	0.0015	0.0000	0.0062	0.0120
AR2 κ	0.3279	***	0.0422	0.0000	0.2451	0.4107
AR2 ϕ_1	1.0072	***	0.0003	0.0000	1.0066	1.0078
AR2 ϕ_2	-0.0113		0.0692	0.8703	-0.1470	0.1244
init	2.8509	***	0.0525	0.0000	2.7480	2.9538
AR1 (2nd) κ	0.4898	***	0.0434	0.0000	0.4048	0.5748
AR1 (2nd) ϕ	0.8628	***	0.0316	0.0000	0.8009	0.9247
seasonal κ	0.0000					
init seasonal 1	0.0075	***	0.0010	0.0000	0.0056	0.0094
init seasonal 2	0.0050	***	0.0011	0.0000	0.0029	0.0071
init seasonal 3	0.0000					
init seasonal 4	0.0000					
init seasonal 5	-0.0125					
β dummy em	0.0184	***	0.0034	0.0000	0.0116	0.0251
β dummy bm	0.0084	*	0.0035	0.0158	0.0016	0.0153
β S&P neg return la..	-0.3428	*	0.1571	0.0292	-0.6508	-0.0348
β WTI Close lag1	-0.2426	***	0.0314	0.0000	-0.3041	-0.1812
Log scale						
AR1 ω	0.0052		0.0036	0.1483	-0.0019	0.0123
AR1 κ	0.1536	***	0.0107	0.0000	0.1327	0.1746
AR1 ϕ	0.9259	***	0.0091	0.0000	0.9081	0.9437
Additional						
Degrees of freedom	7.4019	***	0.7310	0.0000	5.9691	8.8347

* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$

Figure 2
Decomposition of VIX level into AR2, AR1, and seasonal component




Figure 3
Time-varying standard deviation of VIX time series

